






## AIFM2 Antibody

<b>Product Code</b>	CSB-PA874794LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q9BRQ8
Immunogen	Recombinant Human Apoptosis-inducing factor 2 protein (110-373AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, WB, IHC, IF; Recommended dilution: WB:1:500-1:2000, IHC:1:20-1:200, IF:1:50-1:200
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Purification Method	Antigen Affinity Purified
Isotype	IgG
Clonality	Polyclonal
Alias	Apoptosis-inducing factor 2 (EC 1) (Apoptosis-inducing factor homologous mitochondrion-associated inducer of death) (Apoptosis-inducing factor-like mitochondrion-associated inducer of death) (p53-responsive gene 3 protein), AIFM2, AMID PRG3
Immunogen Species	Homo sapiens (Human)
Research Area	Cell Biology
Target Names	AIFM2
Image	IHC image of CSR-PA8747041 A01HH diluted at



IHC image of CSB-PA874794LA01HU diluted at 1:100 and staining in paraffin-embedded human colon cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.




## **CUSABIO TECHNOLOGY LLC**

🕜 Tel: +1-301-363-4651 🛛 Email: cusabio@cusabio.com 🕒 Website: www.cusabio.com 🌘








Western Blot

Positive WB detected in: A549 whole cell lysate

All lanes: AIFM2 antibody at 1:1000

Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 41, 37 kDa Observed band size: 41 kDa



Immunofluorescence staining of HepG2 cells with CSB-PA874794LA01HU at 1:50, counterstained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).