






## PDP1 Antibody

<b>Product Code</b>	CSB-PA878946LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q9P0J1
Immunogen	Recombinant Human [Pyruvate dehydrogenase [acetyl-transferring]]- phosphatase 1, mitochondrial protein (1-154AA)
Raised In	Rabbit
<b>Species Reactivity</b>	Human
<b>Tested Applications</b>	ELISA, WB, IHC; Recommended dilution: WB:1:500-1:5000, IHC:1:500-1:1000
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
<b>Purification Method</b>	>95%, Protein G purified
Isotype	IgG
Clonality	Polyclonal
Alias	[Pyruvate dehydrogenase [acetyl-transferring]]-phosphatase 1, mitochondrial (PDP 1) (EC 3.1.3.43) (Protein phosphatase 2C) (Pyruvate dehydrogenase phosphatase catalytic subunit 1) (PDPC 1), PDP1, PDP PPM2C
Immunogen Species	Homo sapiens (Human)
Research Area	Tags & Cell Markers
Target Names	PDP1

**Image** 



Western Blot

Positive WB detected in: SH-SY5Y whole cell

All lanes: PDP1 antibody at 3.6µg/ml

Goat polyclonal to rabbit IgG at 1/50000 dilution


Predicted band size: 62, 64 kDa Observed band size: 62 kDa



## **CUSABIO TECHNOLOGY LLC**







IHC image of CSB-PA878946LA01HU diluted at 1:600 and staining in paraffin-embedded human testis tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.