





# ZIM2 Antibody

|                            |                                                                                  |
|----------------------------|----------------------------------------------------------------------------------|
| <b>Product Code</b>        | CSB-PA885781LA01HU                                                               |
| <b>Storage</b>             | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.                    |
| <b>Uniprot No.</b>         | Q9NZV7                                                                           |
| <b>Immunogen</b>           | Recombinant Human Zinc finger imprinted 2 protein (1-124AA)                      |
| <b>Raised In</b>           | Rabbit                                                                           |
| <b>Species Reactivity</b>  | Human                                                                            |
| <b>Tested Applications</b> | ELISA, IHC; Recommended dilution: IHC:1:200-1:500                                |
| <b>Form</b>                | Liquid                                                                           |
| <b>Conjugate</b>           | Non-conjugated                                                                   |
| <b>Storage Buffer</b>      | Preservative: 0.03% Proclin 300<br>Constituents: 50% Glycerol, 0.01M PBS, pH 7.4 |
| <b>Purification Method</b> | >95%, Protein G purified                                                         |
| <b>Isotype</b>             | IgG                                                                              |
| <b>Clonality</b>           | Polyclonal                                                                       |
| <b>Alias</b>               | Zinc finger imprinted 2 (Zinc finger protein 656), ZIM2, ZNF656                  |
| <b>Immunogen Species</b>   | Homo sapiens (Human)                                                             |
| <b>Research Area</b>       | Epigenetics and Nuclear Signaling                                                |
| <b>Target Names</b>        | ZIM2                                                                             |

## Image



IHC image of CSB-PA885781LA01HU diluted at 1:300 and staining in paraffin-embedded human brain tissue performed on a Leica Bond<sup>TM</sup> system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC image of CSB-PA885781LA01HU diluted at 1:300 and staining in paraffin-embedded human testis tissue performed on a Leica Bond<sup>TM</sup> system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

