





# HIST1H4A Antibody

|                            |                                                                 |
|----------------------------|-----------------------------------------------------------------|
| <b>Product Code</b>        | CSB-PA951904                                                    |
| <b>Storage</b>             | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.   |
| <b>Uniprot No.</b>         | P62805                                                          |
| <b>Immunogen</b>           | Fusion protein of Human HIST4H4                                 |
| <b>Raised In</b>           | Rabbit                                                          |
| <b>Species Reactivity</b>  | Human,Mouse,Rat                                                 |
| <b>Tested Applications</b> | ELISA,WB,IHC;ELISA:1:2000-1:5000,WB:1:500-1:2000,IHC:1:50-1:200 |
| <b>Form</b>                | Liquid                                                          |
| <b>Conjugate</b>           | Non-conjugated                                                  |
| <b>Storage Buffer</b>      | -20°C, pH7.4 PBS, 0.05% NaN <sub>3</sub> , 40% Glycerol         |
| <b>Purification Method</b> | Antigen affinity purification                                   |
| <b>Isotype</b>             | IgG                                                             |
| <b>Immunogen Species</b>   | Homo sapiens (Human)                                            |
| <b>Target Names</b>        | HIST1H4A                                                        |


## Image



The image on the left is immunohistochemistry of paraffin-embedded Human colon cancer tissue using CSB-PA951904(HIST4H4 Antibody) at dilution 1/20, on the right is treated with fusion protein. (Original magnification: ×200)



The image on the left is immunohistochemistry of paraffin-embedded Human liver cancer tissue using CSB-PA951904(HIST4H4 Antibody) at dilution 1/20, on the right is treated with fusion protein. (Original magnification: ×200)



Gel: 10%SDS-PAGE, Lysate: 40  $\mu$ g, Lane 1-6:  
LoVo cells, NIH/3T3 cells, 293T cells, mouse  
thymus tissue, Hela cells, mouse pancreas  
tissue, Primary antibody: CSB-  
PA951904(HIST4H4 Antibody) at dilution 1/250,  
Secondary antibody: Goat anti rabbit IgG at  
1/8000 dilution, Exposure time: 10 seconds