




Recombinant Human NADH dehydrogenase [ubiquinone] flavoprotein 2, mitochondrial (NDUFV2), partial

Product Code	CSB-RP053854h
Relevance	Core subunit of the mitochondrial mbrane respiratory chain NADH dehydrogenase (Complex I) that is believed to belong to the minimal assbly required for catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone .
Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
Uniprot No.	P19404
Alias	NADH-ubiquinone oxidoreductase 24 kDa subunit
Product Type	Recombinant Protein
Immunogen Species	Homo sapiens (Human)
Purity	Greater than 90% as determined by SDS-PAGE.
Sequence	GGALFVHRDTPENNPDTPTDFPTPENYKRIEAIKKNYPEGHKAAAVLPVLDLAQ RQNGWLPISAMNKVAEVLQVPPMRVYEVAFTYTMYNRKPVGKYHIQVCTTTP CMLRNSDSILEAIQKKLGIKVGETTPDKLFTLIEVECLGACVNAPMVQINDNYE DLTAKDIEEIIDELKAGKIPKPGPRSGRFSCEPAGGLTSLTEPPKGPFGVQAG L
Lead Time	Delivery time may differ from different purchasing way or location, please kindly consult your local distributors for specific delivery time.
Research Area	Transport
Source	E.coli
Gene Names	NDUFV2
Expression Region	35-249aa
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	N-terminal GST-tagged
Mol. Weight	50.6kDa
Protein Description	Partial
Image	



(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.