



## Recombinant Human Putative inactive neutral ceramidase B (ASAH2B)

<b>Product Code</b>	CSB-YP002171HU
Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
Uniprot No.	P0C7U1
Alias	ASAH2-like protein; Putative inactive N-acylsphingosine amidohydrolase 2BPutative inactive non-lysosomal ceramidase B
Product Type	Recombinant Protein
Immunogen Species	Homo sapiens (Human)
Purity	Greater than 90% as determined by SDS-PAGE.
Sequence	MRQHRQFMDRTHYLLTFSSSETLLRLLLRIVDRAPKGRTFGDVLQPAKPEYRV GEVAEVIFVGANPKNSVQNQTHQTFLTVEKYEATSTSWQIVCNDASWETRFY WHKGLLGLSNATVEWHIPDTAQPGIYRIRYFGHNRKQDILKPAVILSFEGTSPA FEVVTI
Lead Time	Delivery time may differ from different purchasing way or location, please kindly consult your local distributors for specific delivery time.
Research Area	Signal Transduction
Source	Yeast
Gene Names	ASAH2B
Protein Names	Recommended name: Putative inactive neutral ceramidase B Alternative name(s): ASAH2-like protein Putative inactive N-acylsphingosine amidohydrolase 2B Putative inactive non-lysosomal ceramidase B
Expression Region	1-165aa
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	N-terminal 6xHis-tagged
Mol. Weight	21.0kDa
<b>Protein Description</b>	Full Length
Image	




## **CUSABIO TECHNOLOGY LLC**

🕜 Tel: +1-301-363-4651 🛛 Email: cusabio@cusabio.com 🕒 Website: www.cusabio.com 💣







(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

## Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL.We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.