



Recombinant Mouse Hyaluronan synthase 2 (Has2), partial

Product Code	CSB-YP010140MO
Relevance	Catalyzes the addition of GlcNAc or GlcUA monosaccharides to the nascent hyaluronan polymer. Therefore, it is essential to hyaluronan synthesis a major component of most Extracellular domain matrices that has a structural role in tissues architectures and regulates cell adhesion, migration and differentiation. This is one of the isozymes catalyzing that reaction and it is particularly responsible for the synthesis of high molecular mass hyaluronan. Required for the transition of endocardial cushion cells into mesenchymal cells, a process crucial for heart development. May also play a role in vasculogenesis. High molecular mass hyaluronan also play a role in early contact inhibition a process which stops cell growth when cells come into contact with each other or the Extracellular domain matrix.
Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
Uniprot No.	P70312
Alias	Hyaluronate synthase 2Hyaluronic acid synthase 2 ;HA synthase 2
Product Type	Recombinant Protein
Immunogen Species	Mus musculus (Mouse)
Purity	Greater than 90% as determined by SDS-PAGE.
Sequence	EHRKMKKSLETPIKLNKTVALCIAAYQEDPDYLRKCLQSVKRLTYPGIKVMVID GNSDDDLMMMDIFSEVMGRDKSATYIWKNNFHEKGPGETEESHKESSQHVT QLVLSNKSICIMQKWGGKREVMYTAFRALGRSVDYVQVCDSDTMLDPASSVE MVKVLEEDPMVGGVGGDVQILNKYDSWISFLSSVRYWMAFNIERACQSYFGC VQCISGPLGMYRNSLLHEFVEDWYNQEFMGNQCSFGDDRHLTNRVLSLGYA TKYTARSKCLTETPIEYLRWLNQQTRWSKSYFREWLYNAMWFHKHHL
Lead Time	Delivery time may differ from different purchasing way or location, please kindly consult your local distributors for specific delivery time.
Research Area	Others
Source	Yeast
Gene Names	Has2
Protein Names	Recommended name: Hyaluronan synthase 2 EC= 2.4.1.212Alternative name(s): Hyaluronate synthase 2 Hyaluronic acid synthase 2 Short name= HA synthase 2
Expression Region	67-374aa
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at



4°C for up to one week.

Tag Info


N-terminal 6xHis-tagged

Mol. Weight

37.9kDa

Protein Description

Partial

Image


(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.