




# Recombinant Rat NADPH oxidase 4 (Nox4), partial

<b>Product Code</b>	CSB-YP015961RA
<b>Relevance</b>	Constitutive NADPH oxidase which generates superoxide intracellularly upon formation of a complex with CYBA/p22phox. Regulates signaling cascades probably through phosphatases inhibition. May function as an oxygen sensor regulating the KCNK3/TASK-1 potassium channel and HIF1A activity. May regulate insulin signaling cascade. May play a role in apoptosis, bone resorption and lipopolysaccharide-mediated activation of NFKB.
<b>Abbreviation</b>	Nox4
<b>Storage</b>	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
<b>Uniprot No.</b>	Q924V1
<b>Alias</b>	Kidney oxidase-1 ;KOX-1Kidney superoxide-producing NADPH oxidase
<b>Product Type</b>	Recombinant Protein
<b>Immunogen Species</b>	Rattus norvegicus (Rat)
<b>Purity</b>	Greater than 90% as determined by SDS-PAGE.
<b>Sequence</b>	GGLLKYQTNLDTHTPPGCISLNRTPSQNMSIADYVSEHFHGSPLGGFSKLEDHY QKTLVKICLEEPKFQAHFPQTWIWISGPLCLYCAERLYRCIRSNKPVITIISVINHP SDVMELRMIKENFKARPGQYIILHCPSVSALENHPFTLTMCPTETKATFGVHFK VVGDWTERFRDLLLPSSQDSEILPFIQSRNYPKLYIDGPFPGSPFEESLNYE
<b>Lead Time</b>	Delivery time may differ from different purchasing way or location, please kindly consult your local distributors for specific delivery time.
<b>Research Area</b>	Others
<b>Source</b>	Yeast
<b>Gene Names</b>	Nox4
<b>Protein Names</b>	Recommended name: NADPH oxidase 4 EC= 1.6.3.-Alternative name(s): Kidney oxidase-1 Short name= KOX-1 Kidney superoxide-producing NADPH oxidase
<b>Expression Region</b>	210-424aa
<b>Notes</b>	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
<b>Tag Info</b>	N-terminal 6xHis-tagged
<b>Mol. Weight</b>	40.6kDa
<b>Protein Description</b>	Partial
<b>Image</b>	



(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

## Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.