




Recombinant Human Complement C4-B (C4B), partial

Product Code	CSB-YP313364HU
Relevance	Non-enzymatic component of the C3 and C5 convertases and thus essential for the propagation of the classical complement pathway. Covalently binds to immunoglobulins and immune complexes and enhances the solubilization of immune aggregates and the clearance of IC through CR1 on erythrocytes. C4A isotype is responsible for effective binding to form amide bonds with immune aggregates or protein antigens, while C4B isotype catalyzes the transacylation of the thioester carbonyl group to form ester bonds with carbohydrate antigens. Derived from proteolytic degradation of complement C4, C4a anaphylatoxin is a mediator of local inflammatory process. It induces the contraction of smooth muscle, increases vascular permeability and causes histamine release from mast cells and basophilic leukocytes.
Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
Uniprot No.	P0C0L5
Alias	Basic complement C4C3 and PZP-like alpha-2-macroglobulin domain-containing protein 3
Product Type	Recombinant Protein
Immunogen Species	Homo sapiens (Human)
Purity	Greater than 90% as determined by SDS-PAGE.
Sequence	EAPKVVEEQESRVHYTVCIWNRNGKVGLSGMAIADVTLLSGFHALRADLEKLTS LSDRYVSHFETEGPHVLLYFDSVPTSRECVGF EAVQEV PVGLVQPASATLYDY YNPERRCSVFY GAPSRSRL LATLCSAEVCQCAEGKCPRQRALERGLQDED GYRMKFACY YPRVEYGFQVKVLREDSRAAFRLFETKITQVLHFTKDVKAAANQ MRNFLVRASCRLRLPEPGKEYLIMGLDGATYDLEGHPQYLLDSNSWIEEMPSE LCRSTRQRAACAQLNDFLQEYGTQGCQV
Lead Time	Delivery time may differ from different purchasing way or location, please kindly consult your local distributors for specific delivery time.
Research Area	Immunology
Source	Yeast
Gene Names	C4B
Protein Names	Recommended name: Complement C4-B Alternative name(s): Basic complement C4 C3 and PZP-like alpha-2-macroglobulin domain-containing protein 3 Cleaved into the following 6 chains: 1. Complement C4 beta chain 2. Complement C4-B alpha chain



Expression Region	1454-1744aa
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	N-terminal 6xHis-tagged
Mol. Weight	35.1kDa
Protein Description	Partial

Image


(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.