



# Recombinant Clostridium botulinum D phage

## Botulinum neurotoxin type D (botD), partial


<b>Product Code</b>	CSB-YP323098CLQ
<b>Relevance</b>	Botulinum toxin acts by inhibiting neurotransmitter release. It binds to peripheral neuronal synapses, is internalized and moves by retrograde transport up the axon into the spinal cord where it can move between postsynaptic and presynaptic neurons. It inhibits neurotransmitter release by acting as a zinc endopeptidase that cleaves the '60-Lys- -Leu-61' bond of synaptobrevins-1 and -2.
<b>Storage</b>	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
<b>Uniprot No.</b>	P19321
<b>Alias</b>	Bontoxilysin-D
<b>Product Type</b>	Recombinant Protein
<b>Immunogen Species</b>	Clostridium botulinum
<b>Purity</b>	Greater than 90% as determined by SDS-PAGE.
<b>Sequence</b>	MTWPVKDFNYSDPVNDNDILYLRI PQNKLITTPVKAFMITQNIWVIPERFSSDTN PSLSKPPRPTSKYQSYDPSYLSSTDEQKDTFLKGIIKLFKRINERDIGKKLINYL VGSPFMGDSSTPEDTFDFTRHTTNIAVEKFENGSWKVNTNIITPSVLIFGPLPNIL DYTASLTQGQQSNPSFEGFGTSLILKVAPEFLLTFSDVTSNQSSAVLGKSIFC MDPVIALMHETHSLHQLYGINIPSDKRIRPQVSEGGFSQDGPVNVQFEELYTFG GLDVEIIPQIERSQLREKALGHYKDIKRLNNINKTIPSSWISNIDKYKKIFSEKYN FDKNTGNFVVNIDKFNSLYSDLTNVMSEVVYSSQYNVKNRTHYFSRHYLPVF ANILDDNIYTIRDFNLTKGFNIENSGQNIERNPALQKLSSSESVVDLFTKVCLR LTK
<b>Lead Time</b>	Delivery time may differ from different purchasing way or location, please kindly consult your local distributors for specific delivery time.
<b>Source</b>	Yeast
<b>Gene Names</b>	botD
<b>Protein Names</b>	Recommended name: Botulinum neurotoxin type D Short name= BoNT/D EC= 3.4.24.69 Alternative name(s): Bontoxilysin-DCleaved into the following 2 chains: 1. Botulinum neurotoxin D light chain 2. Botulinum neurotoxin D heavy chain
<b>Expression Region</b>	1-442aa
<b>Notes</b>	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
<b>Tag Info</b>	N-terminal 6xHis-tagged



**Mol. Weight** 52.5kDa

**Protein Description** Partial

**Image**



(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

**Reconstitution**

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.