



Recombinant Clostridium botulinum Botulinum neurotoxin type F (botF), partial

Product Code	CSB-YP341192CLQ
Relevance	Botulinum toxin acts by inhibiting neurotransmitter release. It binds to peripheral neuronal synapses, is internalized and moves by retrograde transport up the axon into the spinal cord where it can move between postsynaptic and presynaptic neurons. It inhibits neurotransmitter release by acting as a zinc endopeptidase that catalyzes the hydrolysis of the '58-Gln- -Lys-59' bond of synaptobrevins-1 and -2.
Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
Uniprot No.	P30996
Alias	Bontoxilysin-F
Product Type	Recombinant Protein
Immunogen Species	Clostridium botulinum
Purity	Greater than 90% as determined by SDS-PAGE.
Sequence	MPVAINSFNYNDPVNDDTILYMQIPYEEKSKKYYKA FEIMRNVWIIPERN TIGTN PSDFDPPASLKN GSSAYYDPNYLTTDAEKDRYLKTTIKL FKRINSNPAGKVLLQ EISYAKPYLGNDHTPID EFSVTRTTSVNIKLSTNVESSMLLNLLVLGAGPDIFE SCCYPVRKLIDPDVVYDPSNYGFGSINIVTFSPEYEYTFNDISGGHNSSTESFIA DPAISLAHELIALHGLYGARGVTYEETIEVKQAPLMIAEKPIRLEEF LTFGGQDL NIITSAMKEKIYNNLLANYEKIATRLSEVNSAPPEYDINEYKDYFQWKYGLDKNA DGSYTVNENKFNEIYKKLYSFTESDLANKFKVKCRNTYFIKYEF LKVPNLLDDD YTVSEGFNIGNLAVNNRGQSIKLNPKIIDSIPDKGLVEKIVKFKCKSVIPRK
Lead Time	Delivery time may differ from different purchasing way or location, please kindly consult your local distributors for specific delivery time.
Research Area	Others
Source	Yeast
Gene Names	botF
Protein Names	Recommended name: Botulinum neurotoxin type F Short name= BoNT/F EC= 3.4.24.69 Alternative name(s): Bontoxilysin-F Cleaved into the following 2 chains: 1. Botulinum neurotoxin F light chain 2. Botulinum neurotoxin F heavy cha
Expression Region	1-436aa
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.

**Tag Info**


N-terminal 6xHis-tagged

Mol. Weight

51.6kDa

Protein Description

Partial

Image

(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.