




Recombinant Mouse Histone deacetylase complex subunit SAP130 (Sap130), partial

| | |
|----------------------------|---|
| Product Code | CSB-YP807359MO |
| Relevance | Acts as a transcriptional repressor. May function in the assembly and/or enzymatic activity of the mSin3A corepressor complex or in mediating interactions between the complex and other regulatory complexes |
| Abbreviation | Sap130 |
| Storage | The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C. |
| Uniprot No. | Q8BIH0 |
| Alias | 130 kDa Sin3-associated polypeptide Sin3-associated polypeptide p130 |
| Product Type | Recombinant Protein |
| Immunogen Species | Mus musculus (Mouse) |
| Purity | Greater than 90% as determined by SDS-PAGE. |
| Sequence | PRKQQHVISTEEGDMMETNSTDDEKSAKSLLVKAEKRKSPPEYIDEEGVRY VPVRPRPPITLLRHYRNPWKAAYHHFQRYSDVRVKEEKKAMLQEIANQKGVS CRAQGWKVHLCAAQLQLTNLEHDVYERLTNLQEGIIIPKKKAATDDDLHRINEL IQGNMQRCKLVMDQISEARDSMLKVLDPKDRVLKLLNKNKGTGKKVSKLKRKEK V |
| Lead Time | Delivery time may differ from different purchasing way or location, please kindly consult your local distributors for specific delivery time. |
| Research Area | Others |
| Source | Yeast |
| Gene Names | Sap130 |
| Protein Names | Recommended name: Histone deacetylase complex subunit SAP130 Alternative name(s): 130 kDa Sin3-associated polypeptide Sin3-associated polypeptide p130 |
| Expression Region | 845-1057aa |
| Notes | Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week. |
| Tag Info | N-terminal 6xHis-tagged |
| Mol. Weight | 26.8kDa |
| Protein Description | Partial |
| Image | |



(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.