

## Olfactory receptor 2A25 rabbit pAb

Cat No.:ES3025

For research use only

## Overview

Product Name Olfactory receptor 2A25 rabbit pAb

Host species Rabbit
Applications WB;IF;ELISA

Species Cross-Reactivity Human; Rat; Mouse;

**Recommended dilutions** Western Blot: 1/500 - 1/2000. Immunofluorescence:

1/200 - 1/1000. ELISA: 1/5000. Not yet tested in

other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human OR2A25. AA

range:241-290

**Specificity** Olfactory receptor 2A25 Polyclonal Antibody detects

endogenous levels of Olfactory receptor 2A25

protein.

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

**Storage** Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Olfactory receptor 2A25

Gene Name OR2A25

Cellular localizationCell membrane; Multi-pass membrane protein.PurificationThe antibody was affinity-purified from rabbit<br/>antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 40kD
Human Gene ID 392138
Human Swiss-Prot Number A4D2G3

Alternative Names OR2A25; OR2A25p; OR2A27; Olfactory receptor

2A25; Olfactory receptor 2A27


**Background** Olfactory receptors interact with odorant molecules

in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of






G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],




Western Blot analysis of various cells using Olfactory receptor 2A25 Polyclonal Antibody




Immunofluorescence analysis of A549 cells, using OR2A25 Antibody. The picture on the right is blocked with the synthesized peptide.







Western blot analysis of lysates from HT-29 cells, using OR2A25 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HepG2 cells using OR2A25 antibody.



23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei, P.R.C