

## Nop132 rabbit pAb

Cat No.: ES6693

For research use only

## Overview

Product Name Nop132 rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA Species Cross-Reactivity Human;Rat;Mouse;

**Recommended dilutions** Western Blot: 1/500 - 1/2000.

Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human NOL8. AA

range:1118-1167

**Specificity** Nop132 Polyclonal Antibody detects endogenous

levels of Nop132 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

**Storage** Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Nucleolar protein 8

Gene Name NOL8

**Cellular localization** Nucleus, nucleolus . Localizes in the

nucleolar-organizing region during ribosome

biogenesis.

**Purification** The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 130kD
Human Gene ID 55035
Human Swiss-Prot Number Q76FK4

Alternative Names NOL8; C9orf34; NOP132; Nucleolar protein 8;

Nucleolar protein Nop132

Background NOL8 binds Ras-related GTP-binding proteins (see


MIM 608267) and plays a role in cell growth

(Sekiguchi et al., 2004 [PubMed




+86-27-59760950 ELKbio@ELKbiotech.com www.elkb






Western Blot analysis of 3T3 cells using Nop132 Polyclonal Antibody diluted at 1:1000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).




3T3

Western Blot analysis of NIH-3T3 cells using Nop132 Polyclonal Antibody diluted at 1:1000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).




Western blot analysis of NOL8 Antibody. The lane on the right is blocked with the NOL8 peptide.



+86-27-59760950







Immunohistochemistryt analysis of paraffin-embedded human brain, using NOL8 Antibody. The lane on the right is blocked with the NOL8 peptide.



+86-27-59760950