

SNX4 rabbit pAb

Cat No.: ES7866

For research use only

Overview

Product Name SNX4 rabbit pAb

Host species Rabbit
Applications WB;IHC

Species Cross-Reactivity Human; Mouse; Rat

Recommended dilutions WB 1:500-2000;IHC-p 1:50-300

Immunogen The antiserum was produced against synthesized

peptide derived from human SNX4. AA

range:191-240

Specificity SNX4 Polyclonal Antibody detects endogenous levels

of SNX4 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Sorting nexin-4

Gene Name SNX4

Cellular localization Early endosome membrane; Peripheral membrane

protein; Cytoplasmic side. Also detected on a juxtanuclear endocytic recycling compartment

(ERC). .

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 60kD
Human Gene ID 8723
Human Swiss-Prot Number 095219

Alternative Names SNX4; Sorting nexin-4

Background This gene encodes a member of the sorting nexin

family. Members of this family contain a phox (PX) domain, which is a phosphoinositide binding

domain, and are involved in intracellular trafficking. This protein associated with the long isoform of the




+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com




leptin receptor and with receptor tyrosine kinases for platelet-derived growth factor, insulin, and epidermal growth factor in cell cultures, but its function is unknown. This protein may form oligomeric complexes with family members. Two transcript variants, one protein coding and the other non-protein coding, have been found for this gene. [provided by RefSeq, Nov 2012],

Western blot analysis of the lysates from HeLa cells using SNX4 antibody.



Immunohistochemical analysis of paraffin-embedded human Squamous cell carcinoma of lung. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).



+86-27-59760950