

## Ezrin (Phospho-Thr566) Antibody

Catalog Number: E011202-1, E011202-2

**Amount:** 50μg/50μl, 100μg/100μl

Swiss-Prot No.: P15311

All Names: Cytovillin, EZRI, VIL2, Villin 2, p81

All Sites: Human: Thr566 or Thr567; Mouse: Thr566 or Thr567; Rat: Thr567

Form of Antibody: Rabbit IgG in phosphate buffered saline (without Mg<sup>2+</sup> and Ca<sup>2+</sup>), pH 7.4, 150mM NaCl,

0.02% sodium azide and 50% glycerol.

Storage/Stability: Store at -20°C/1 year

Immunogen: The antiserum was produced against synthesized phosphopeptide derived from human

Ezrin around the phosphorylation site of threonine 566 (Y-K-T<sup>P</sup>-L-R).

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using

epitope-specific phosphopeptide. The antibody against non-phosphopeptide was removed by chromatogramphy using non-phosphopeptide corresponding to the phosphorylation site.

**Specificity/Sensitivity:** Ezrin (phospho-Tyr566) antibody detects endogenous levels of Ezrin only when

phosphorylated at threonine 566.


**Reactivity:** Human, Mouse, Rat **Applications:** WB: 1:500~1:1000

References: Zhou R, et al. (2005) J Cell Sci: 118 (Pt 19): 4381-91

Shiue H, et al.(2005) J Biol Chem: 280(2): 1688-95

Zhao H, et al. (2004) Proc Natl Acad Sci U S A: 101(25): 9485-90

Fievet BT, et al. (2004) J Cell Biol: 164(5): 653-9



Western blot analysis of extracts from A431 cells, untreated or EGF-treated (200ng/ml, 30min) using Ezrin (Ab-566) antibody (E021200, Lane 1 and 2) and Ezrin (phospho-Thr566) antibody

(E011202, Lane 3 and 4).