

Smad2 (Phospho-Ser467) Antibody

Catalog Number: E011322-1, E011322-2

Amount: 50μg/50μl, 100μg/100μl

Swiss-Prot No.: Q15796

Form of Antibody: Rabbit IgG in phosphate buffered saline (without Mg²⁺ and Ca²⁺), pH 7.4, 150mM NaCl,

0.02% sodium azide and 50% glycerol.

Storage/Stability: Store at -20 °C/1 year

Immunogen: The antiserum was produced against synthesized phosphopeptide derived from human

Smad2 around the phosphorylation site of serine 467 (C-S-S-M-S^P).

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using

epitope-specific phosphopeptide. The antibody against non-phosphopeptide was removed by chromatography using non-phosphopeptide corresponding to the phosphorylation site.


Specificity/Sensitivity: Smad2 (Phospho-Ser467) Antibody detects endogenous levels of Smad2 only when

phosphorylated at serine 467.

Reactivity: Human, Mouse, Rat **Applications:** WB: 1:500~1:1000

References: Sang Gyun Kim et al. (2004) Mol Biol Cell. February; 15(2): 420–434.

Rebecca A. Randall et al. (2004) Mol Cell Biol. February; 24(3): 1106–1121. Kimberly A Brown et al. (2004) Breast Cancer Res.; 6(3): R215–R231.



Western blot analysis of extract from 293 cells, untreated or treated with PMA (200nM, 30min), using Smad2 (Ab-467) Antibody (E021322, Lane 1 and 2) and Smad2 (Phospho-Ser467) Antibody (E011322, Lane 3 and 4).