

FLT1 Mouse Monoclonal Antibody

Background:

Fms-related tyrosine kinase 1 (vascular endothelial growth factor/vascular permeability factor receptor), also known as FLT1 or VEGFR-1. It is a member of the vascular endothelial growth factor receptor (VEGFR) family. VEGFR family members are receptor tyrosine kinases (RTKs) which contain an extracellular ligand-binding region with seven immunoglobulin (Ig)-like domains, a transmembrane segment, and a tyrosine kinase (TK) domain within the cytoplasmic domain. This protein binds to VEGFR-A, VEGFR-B and placental growth factor and plays an important role in angiogenesis and vasculogenesis. Expression of this receptor is found in vascular endothelial cells, placental trophoblast cells and peripheral blood monocytes. Multiple transcript variants encoding different isoforms have been found for this gene. Isoforms include a full-length transmembrane receptor isoform and shortened, soluble isoforms. The soluble isoforms are associated with the onset of pre-eclampsia.

Catalog Number: E10-20387

Amount: 100μg/100μl

Clone Number: 3D10

Species: Mouse IgG1
MW: 100kDa
Aliases: FLT; VEGFR1

Entrez Gene: 2321

Immunogen: Purified recombinant extracellular fragment of human FLT1 fused with hlgGFc tag expressed in

HEK293 cells.

Storage: Store at 4 °20 for Cong term storage, store at


Formulation: Ascitic fluid containing 0.03% sodium azide.

Species Reactivities: Human

Tested Applications: WB, ,ELISA. Not yet tested in other applications. Determining optimal working dilutions by

titration test.

Application notes: WB. 1/500 - 1/2000. ELISA. Propose dilution 1/10000.



rigure 1. Western blot analysis using the finduse mab against extracellular domain of numarither t