

CDH1 Mouse Monoclonal

Antibody

Background:

E-Cadherin is a 120 kDa transmembrane glycoprotein that is localized in the adherens junctions of epithelial cells. There, it interacts with the cytoskeleton through the associated cytoplasmic catenin proteins. In addition to being a calcium-dependent adhesion molecule, E-Cadherin is also a critical regulator of epithelial junction formation. Its association with catenins is necessary for cell-cell adhesion. These E-cadherin/catenin complexes associate with corical actin bundles at both the zonula adherens and the lateral adhesion plaques. Tyrosine phosphorylation can disrupt these complexes, leading to changes in cell adhesion properties. E-Cadherin expression is often down-regulated in highly invasive, poorly differentiated carcinomas. Increased expression of E-Cadherin in these cells reduces invasiveness. Thus, loss of expression or function of E-Cadherin appears to be an important step in tumorigenic progression. Tissue specificity: Non-neural epithelial tissues.

Catalog Number: E10-30127

Amount: 100μg/100μl

Clone Number: 7A2

Species: Mouse IgG1 **MW:** 135kDa

Aliases: UVO; CDHE; ECAD; LCAM; Arc-1; CD324; CDH1

Entrez Gene: 999

Immunogen: Purified recombinant fragment of human CDH1 expressed in E. Coli.

Storage: Store at 4° C, for long term storage, store at -20° C.

Formulation: Ascitic fluid containing 0.03% sodium azide.

Species Reactivities: Human; Mouse; Monkey

Tested Applications: WB, ELISA. Not yet tested in other applications.

Application notes: WB: 1/500 - 1/2000,

ELISA: Propose dilution 1/10000.




Figure 1: Western blot analysis using CDH1 mouse mAb against LNCAP (1), A431 (2), DU145 (3), PC-3 (4), PC-12 (5) and T47D(6) cell lysate.