



BAD (Cleaved-Asp71) Antibody

E11-0102L

Catalog Number: E11-0102L

Amount: 100 μ g/100 μ l

Swiss-Prot No. : Q92934

All Names: BBC2, BCL2L8, BCL-X/BCL-2 binding protein, BCL2 ANTAGONIST OF CELL DEATH
BCL2-antagonist of cell death protein, BCL2-binding component 6, BCL2-binding protein

All Sites: Mouse: Asp71; Rat: Asp71

Form of Antibody: Rabbit IgG in phosphate buffered saline (without Mg²⁺ and Ca²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

Storage/Stability: Store at -20°C/1 year

Immunogen: The antiserum was produced against synthesized peptide derived from mouse BAD.

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.


Specificity/Sensitivity: BAD (Cleaved-Asp71) antibody detects endogenous levels of fragment of activated BAD resulting from cleavage adjacent to Asp71.

Reactivity: Mouse, Rat

Applications: WB: 1:300~1000 IHC: 1:50~100


ELISA: 1:20000

References: W. Steven Donaldson, Bulletin of Canadian Petroleum Geology, Dec 1998; 46: 576 - 598.
DALE TSHUDY, Journal of Paleontology, Sep 2005; 79: 961 - 968.
Alexandre David, J. Cell Sci., Jan 2007; 120: 353 - 365.
Peng Jiang, Mol. Cell. Biol., Dec 2006; 26: 9071 - 9082.



CI-Peptide - +

Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using BAD (Cleaved-Asp71) antibody.



Western blot analysis of extracts from 293 cells, treated with Etoposide (25uM, 60mins), using BAD (Cleaved-Asp71) antibody.

For Research Use Only