

Catalog Number:E11-0493A
Concentration: 1mg/ml

Swiss-Prot No.: P06213

Other Names: CD220 antigen, EC 2.7.10.1, IR, insulin receptor, kinase InsR

All Sites: Human: Tyr1355; Mouse: Tyr1345; Rat: Tyr1356

Storage/Stability: Store at -20°C/1 year

Form of Antibody: Rabbit IgG in phosphate buffered saline (without Mg²⁺ and Ca²⁺), pH 7.4, 150mM NaCl, 0.02%

sodium azide and 50% glycerol.

 $\textbf{Immunogen:} \ \textbf{The antiserum was produced against synthesized phosphopeptide derived from human IR around the} \\$

phosphorylation site of tyrosine 1355 (R-S-Y^P-E-E).

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific phosphopeptide. The antibody against non-phosphopeptide was removed by chromatography using non-phosphopeptide corresponding to the phosphorylation site.

Specificity: IR (Phospho-Tyr1355) antibody detects endogenous levels of IR only when phosphorylated at tyrosine 1355.


Reactivity: Human, Mouse, Rat

References:

Laura Sciacca, Endocrinology, Jun 2003; 144: 2650.

Mark Westcott, Asia Pacific Journal of Human Resources, Aug 2003; 41: 172 - 189.

Adam Denley, Mol. Endocrinol., Oct 2004; 18: 2502 - 2512. Rita Slaaby, J. Biol. Chem., Sep 2006; 281: 25869 - 25874. -----



with Heat shock, using IR (Phospho-Tyr1355) antibody.