



BCL-2 (Ab-69) Antibody

E11-0774B**Catalog Number:** E11-0774B**Concentration:** 1mg/ml**Swiss-Prot No.:** P10415**Other Names:** Apoptosis regulator Bcl-2, BCL2**Storage/Stability:** Store at -20°C/1 year**Form of Antibody:** Rabbit IgG in phosphate buffered saline (without Mg^{2+} and Ca^{2+}), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.**Immunogen:** The antiserum was produced against synthesized non-phosphopeptide derived from human BCL-2 around the phosphorylation site of threonine 69 (A-R-T^P-S-P).**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using

epitope-specific immunogen.


Specificity: BCL-2 (Ab-69) antibody detects endogenous levels of total BCL-2 protein.**Reactivity:** Human, Mouse, Rat**Applications:** WB: 1:500~1:1000 IHC: 1:50~1:100
IF: 1:100~1:500 ELISA: 1:10000**References:**

H. Kurosawa, Am J Physiol Gastrointest Liver Physiol, Jun 1997; 272: 1587.

JR Park, Blood, Aug 1995; 86: 868 - 876.


A Plettenberg, Am. J. Pathol., Mar 1995; 146: 651.

JM Ben-Ezra, Am. J. Pathol., Nov 1994; 145: 1036.




Peptide - +

Immunohistochemistry analysis of paraffin-embedded human colon carcinoma tissue using BCL-2 (Ab-69) antibody.



Western blot analysis of extracts from K562 cells, using BCL-2 (Ab-69) antibody.

For Research Use Only



Peptide

-

+

Immunofluorescence analysis of HepG2 cells, using
BCL-2 (Ab-69) antibody.