

EDG4 Antibody

Catalog Number: E11-085G

Concentration: 1mg/ml,100ug/100uL

Swiss-Prot No.: Q9HBW0

Other Names: EDG-4; endothelial differentiation, lysophosphatidic acid G-protein-coupled receptor, 4; G protein-coupled receptor; LPA receptor 2; LPA receptor EDG4; LPA-2; LPA2; LPAR2; lysophosphatidic acid receptor Edg-4; lysophosphatidic acid receptor EDG4 Storage/Stability: Store at -20 °C/1 year Form of Antibody: Rabbit IgG in phosphate buffered saline (without Mg²⁺ and Ca²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

Immunogen: The antiserum was produced against synthesized peptide derived from internal of human

EDG4.

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.


Specificity: EDG4 antibody detects endogenous levels of

total EDG4 protein. Reactivity: Human


Applications: WB: 1:500~1:1000 IF: 1:100~1:500

ELISA: 1:10000 References:

An S., J. Biol. Chem. 273:7906-7910(1998). Bandoh K., FEBS Lett. 478:159-165(2000). Grimwood J., Nature 428:529-535(2004).



Western blot analysis of extracts from COLO205 cells, using EDG4 antibody.



Immunofluorescence analysis of LOVO cells, using EDG4 antibody.