



c-Met (Ab-1003) Antibody

E11-0871B**Catalog Number:** E11-0871B**Concentration:** 1mg/ml**Swiss-Prot No.:** P08581

Other Names: EC 2.7.10.1, HGF receptor, HGF-SF receptor, Hepatocyte growth factor receptor precursor, Met proto-oncogene tyrosine kinase, c-met, kinase EC 2.7.10.1, HGF receptor, HGF-SF receptor, Hepatocyte growth factor receptor precursor, Met proto-oncogene tyrosine kinase, c-met, kinase Met

Storage/Stability: Store at -20°C/1 year

Form of Antibody: Rabbit IgG in phosphate buffered saline (without Mg^{2+} and Ca^{2+}), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

Immunogen: The antiserum was produced against synthesized non-phosphopeptide derived from human c-Met around the phosphorylation site of tyrosine 1003 (V-D-Y^P-R-A).

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Specificity: c-Met (Ab-1003) antibody detects endogenous levels of total c-Met protein.

Reactivity: Human, Mouse, Rat


Applications: WB: 1:500~1:1000 IF: 1:100~1:500 ELISA: 1:1000

References:


Hillier L.W., Nature 424:157-164(2003).

Wang D., J. Biol. Chem. 277:36216-36222(2002).

Giordano S., Nat. Cell Biol. 4:720-724(2002).



Western blot analysis of extracts from HepG2 cells, using c-Met (Ab-1003) antibody.



Immunofluorescence analysis of A549 cells, using c-Met (Ab-1003) antibody.

For Research Use Only