

PHLA2 Antibody

Catalog Number: E11-11168 C

Concentration: 1mg/ml Swiss-Prot No.: Q53GA4

Other Names: Imprinted in placenta and liver protein; Tumor-suppressing subchromosomal transferable fragment candidate gene 3 protein; Tumor-suppressing STF cDNA 3 protein; Beckwith-Wiedemann syndrome chromosomal region 1 candidate gene C protein; p17-Beckwith-Wiedemann region 1 C; p17-BWR1C

Storage/Stability: Store at -20 °C/1 year

Form of Antibody: Rabbit IgG in phosphate buffered saline (without Mg²⁺ and Ca²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

Immunogen: The antiserum was produced against

synthesized peptide derived from internal of human PHLA2.

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Specificity: PHLA2 antibody detects endogenous levels of total PHLA2 protein.

Reactivity: Human (Identities = 100%, Positives = 100%) **Applications:** IHC: 1:50~1:100 ELISA: 1:20000

References:

Hu R.-J., Genomics 46:9-17(1997).

Qian N., Hum. Mol. Genet. 6:2021-2029(1997).

 $\label{eq:constraints} \text{Schwienbacher} \quad \text{C.,} \quad \text{Proc.} \quad \text{Natl.} \quad \text{Acad.} \quad \text{Sci.} \quad \text{U.S.A.}$

95:3873-3878(1998).

Peptide - +

Immunohistochemistry analysis of paraffin-embedded human placenta tissue, using PHLA2 antibody.