



BCL10 Antibody

Catalog Number: E11-13030 C

Concentration: 1mg/ml Swiss-Prot No.: O95999

Other Names: B-cell CLL/lymphoma 10, B-cell lymphoma/leukemia 10, BCL10, CARD- containing molecule enhancing NF-kappa-B, CARD-like apoptotic protein, CED- 3/ICH-1 prodomain homologous E10-like regulator, CIPER, CLAP, Cellular homolog of vCARMEN, Cellular-E10, Mammalian CARD-containing adapter molecule E10, c-E10, cCARMEN, hCLAP, mE10

Storage/Stability: Store at -20°C/1 year

Form of Antibody: Rabbit IgG in phosphate buffered saline (without Mg²⁺ and Ca²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

Immunogen: The antiserum was produced against synthesized peptide derived from internal of human

BCL10.

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using

epitope-specific immunogen.

Specificity: BCL10 antibody detects endogenous levels

of total BCL10 protein.

Reactivity: Human (Identities = 100%, Positives = 100%);

Mouse (Identities = 85%, Positives = 85%);

Rat (Identities = 100%, Positives = 100%)

ELISA: 1:40000
References:

Willis T.G., Cell 96:35-45(1999).

Koseki T., J. Biol. Chem. 274:9955-9961(1999).

Thome M., J. Biol. Chem. 274:9962-9968(1999).

Peptide - +

Immunohistochemistry analysis of paraffin-embedded human colon carcinoma tissue using BCL10 antibody.

K562 K562
-- 117
-- 85
-- 48
BCL10-- -- 34
-- 26
-- 19
(kD)
Peptide - +

Western blot analysis of extracts from K562 cells, using BCL10 antibody.