




NF-κB p65 (Phospho-Ser529) Antibody

E11-7162A

- Catalog Number:** E11-7162A
- Amount:** 100µg/100µl
- Swiss-Prot No.:** Q04206
- All Names:** NF-KB p65, NFKB3, Nuclear factor NF-kappa-B p65 subunit, RELA, TF65, Transcription factor p65, p65
- All Sites:** Human: Ser529; Mouse: Ser529
- Form of Antibody:** Rabbit IgG in phosphate buffered saline (without Mg^{2+} and Ca^{2+}), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
- Storage/Stability:** Store at -20 °C/1 year
- Immunogen:** The antiserum was produced against synthesized non-phosphopeptide derived from human NF-κB p65 around the phosphorylation site of serine 529 (L-L-S^P-G-D).
- Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific phosphopeptide. The antibody against non-phosphopeptide was removed by chromatography using non-phosphopeptide corresponding to the phosphorylation site.
- Specificity/Sensitivity:** NF-κB p65 (Ab-529) antibody detects endogenous levels of NF-κB p65 protein only when phosphorylated at serine 529.
- Reactivity:** Human, Mouse
- Applications:** WB: 1:500~1:1000 IHC: 1:50~1:100
ELISA: 1:10000
- References:** Xu C, et al (2005) Oncogene:24(28): 4486-95.
McNulty SE, et al. (2004) Pigment Cell Res Apr; 17(2): 173-80.
Madrid LV, et al. (2001) J Biol Chem: 276(22): 18934-40.
Wang D, et al. (2000) J Biol Chem : 275(42): 32592-7.




P-Peptide

-

+

Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using NF-κB p65 (phospho-Ser529) antibody.



Western blot analysis of extracts from MDA-MB-231 cells, untreated or treated with TNF-α (20ng/ml, 10min) using NF-κB p65 (Ab-529) antibody (Line 1 and 2) and NF-κB p65 (phospho-Ser529) antibody (Line 3 and 4).

For Research Use Only