



| | | |
|--|--|--|
|  | <h2>Cleaved-Integrin α5 LC (E895) Polyclonal Antibody</h2> | |
|--|--|--|

| | |
|---------------------------------|---|
| Swiss-Prot No.: | P08648 |
| Molecular Weight: | 114.536 |
| Applications: | WB,ELISA |
| Reactivity: | Human |
| Purification: | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Immunogen: | Synthesized peptide derived from the C-terminal region of human Integrin α 5 LC. at AA range: 850-930 |
| Specificity/Sensitivity: | Cleaved-Integrin α 5 LC (E895) Polyclonal Antibody detects endogenous levels of fragment of activated Integrin α 5 LC protein resulting from cleavage adjacent to E895. |
| Other Names: | ITGA5; FNRA; Integrin alpha-5; CD49 antigen-like family member E; Fibronectin receptor subunit alpha; Integrin alpha-F; VLA-5; CD antigen CD49e |
| Storage/Stability: | -20°C/1 year |
| Source: | Rabbit |
| Form of Antibody: | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Protein Concentration: | 1 mg/ml |

For Research Use Only

| | |
|-------------------|---|
| |  <p>Western blot analysis showing a single band at approximately 19 kD. Molecular weight markers are indicated on the left: 293, 117, 85, 48, 34, 26, and 19 kD.</p> |
| | Western Blot analysis of various cells using Cleaved-Integrin $\alpha 5$ LC (E895) Polyclonal Antibody |
| Gene Name: | ITGA5 |
| Dilution: | Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications. |