




## Met (phospho Tyr1356) Polyclonal Antibody

<b>Swiss-Prot No.:</b>	P08581
<b>Molecular Weight:</b>	155.527
<b>Applications:</b>	WB,ELISA
<b>Reactivity:</b>	Human,Mouse,Rat
<b>Purification:</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Immunogen:</b>	Synthesized peptide derived from human Met around the phosphorylation site of Y1356.
<b>Specificity/Sensitivity:</b>	Phospho-Met (Y1356) Polyclonal Antibody detects endogenous levels of Met protein only when phosphorylated at Y1356.
<b>Other Names:</b>	MET; Hepatocyte growth factor receptor; HGF receptor; HGF/SF receptor; Proto-oncogene c-Met; Scatter factor receptor; SF receptor; Tyrosine-protein kinase Met
<b>Storage/Stability:</b>	-20°C/1 year
<b>Source:</b>	Rabbit
<b>Form of Antibody:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Protein Concentration:</b>	1 mg/ml

	
	Western Blot analysis of 3T3 cells using Phospho-Met (Y1356) Polyclonal Antibody
Gene Name:	MET
Dilution:	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.