


	<p>p27 (phospho Thr198) Polyclonal Antibody</p>	
--	---	--

<b>Swiss-Prot No.:</b>	P46527
<b>Molecular Weight:</b>	22.073
<b>Applications:</b>	WB,ELISA
<b>Reactivity:</b>	Human
<b>Purification:</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Immunogen:</b>	Synthesized peptide derived from human p27 around the phosphorylation site of T198.
<b>Specificity/Sensitivity:</b>	Phospho-p27 (T198) Polyclonal Antibody detects endogenous levels of p27 protein only when phosphorylated at T198.
<b>Other Names:</b>	CDKN1B; KIP1; Cyclin-dependent kinase inhibitor 1B; Cyclin-dependent kinase inhibitor p27; p27Kip1
<b>Storage/Stability:</b>	-20°C/1 year
<b>Source:</b>	Rabbit
<b>Form of Antibody:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Protein Concentration:</b>	1 mg/ml

	 <p>Western blot analysis of VEC cells using Phospho-p27 (T198) Polyclonal Antibody. The blot shows a single band at approximately 25 kDa, labeled p-p27 (T198). Molecular weight markers are indicated on the left: 170, 130, 100, 70, 55, 40, 35, 25, and 15 kDa.</p>
	Western Blot analysis of VEC cells using Phospho-p27 (T198) Polyclonal Antibody
<b>Gene Name:</b>	CDKN1B
<b>Dilution:</b>	Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.