
	<h2>Smad2/3 (phospho Thr8) Polyclonal Antibody</h2>	
--	---	--

<b>Swiss-Prot No.:</b>	Q15796/P84022
<b>Molecular Weight:</b>	48.081
<b>Applications:</b>	WB,ELISA
<b>Reactivity:</b>	Human,Mouse,Rat
<b>Purification:</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Immunogen:</b>	Synthesized peptide derived from human Smad2/3 around the phosphorylation site of T8.
<b>Specificity/Sensitivity:</b>	Phospho-Smad2/3 (T8) Polyclonal Antibody detects endogenous levels of Smad2/3 protein only when phosphorylated at T8.
<b>Other Names:</b>	SMAD2; MADH2; MADR2; Mothers against decapentaplegic homolog 2; MAD homolog 2; Mothers against DPP homolog 2; JV18-1; Mad-related protein 2; hMAD-2; SMAD family member 2; SMAD 2; Smad2; hSMAD2; SMAD3; MADH3; Mothers against decapentaplegic
<b>Storage/Stability:</b>	-20°C/1 year
<b>Source:</b>	Rabbit
<b>Form of Antibody:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

**For Research Use Only**

<b>Protein Concentration:</b>	1 mg/ml
	 <p>Western blot analysis of VEC cells showing p-Smad2/3 (T8) phosphorylation. The blot shows a single band at approximately 66 kDa. Molecular weight markers are indicated on the left: 170, 130, 100, 70, 55, 45, 35, 25, and 15 kDa.</p>
	Western Blot analysis of various cells using Phospho-Smad2/3 (T8) Polyclonal Antibody
<b>Gene Name:</b>	SMAD2/SMAD3
<b>Dilution:</b>	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.