



| | | |
|--|---|--|
|  | <h2 style="text-align: center;">Flt-3 (phospho Tyr842) Polyclonal Antibody</h2> | |
|--|---|--|

| | |
|---------------------------------|---|
| Swiss-Prot No.: | P36888 |
| Molecular Weight: | 112.804 |
| Applications: | WB,ELISA |
| Reactivity: | Human,Mouse |
| Purification: | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Immunogen: | Synthesized peptide derived from human Flt-3 around the phosphorylation site of Y842. |
| Specificity/Sensitivity: | Phospho-Flt-3 (Y842) Polyclonal Antibody detects endogenous levels of Flt-3 protein only when phosphorylated at Y842. |
| Other Names: | FLT3; CD135; FLK2; STK1; Receptor-type tyrosine-protein kinase FLT3; FL cytokine receptor; Fetal liver kinase-2; FLK-2; Fms-like tyrosine kinase 3; FLT-3; Stem cell tyrosine kinase 1; STK-1; CD antigen CD135 |
| Storage/Stability: | -20°C/1 year |
| Source: | Rabbit |
| Form of Antibody: | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Protein Concentration: | 1 mg/ml |

For Research Use Only

| | |
|-------------------|--|
| |  <p>293T-UV</p> <p>170— 130— 100— 70— 55— 40— 35— 25— 15—</p> <p>p-Flt-3 (Y842)</p> |
| | Western Blot analysis of 293T-UV cells using Phospho-Flt-3 (Y842) Polyclonal Antibody |
| Gene Name: | FLT3 |
| Dilution: | Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. |