



## PAKa/β/γ (phospho Thr423/402/421) Polyclonal Antibody

|                                 |  |
|---------------------------------|--|
| <b>Swiss-Prot No.:</b>          | Q13153/Q13177/075914   |
| <b>Applications:</b>            | WB,IHC,ELISA   |
| <b>Reactivity:</b>              | Human,Mouse,Rat  |
| <b>Purification:</b>            | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  |
| <b>Immunogen:</b>               | Synthesized peptide derived from human PAKa/β/γ around the phosphorylation site of T423/402/421.   |
| <b>Specificity/Sensitivity:</b> | Phospho-PAKa/β/γ (T423/402/421) Polyclonal Antibody detects endogenous levels of PAKa/β/γ protein only when phosphorylated at T423/402/421.  |
| <b>Other Names:</b>             | PAK1; Serine/threonine-protein kinase PAK 1; Alpha-PAK; p21-activated kinase 1; PAK-1; p65-PAK; PAK2; Serine/threonine-protein kinase PAK 2; Gamma-PAK; PAK65; S6/H4 kinase; p21-activated kinase 2; PAK-2; p58; PAK3; OPHN3; Serine/threonine-p |
| <b>Storage/Stability:</b>       | -20°C/1 year   |
| <b>Source:</b>                  | Rabbit   |
| <b>Form of Antibody:</b>        | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |

|                               |   |
|-------------------------------|---|
| <b>Protein Concentration:</b> | 1 mg/ml   |
| <b>Gene Name:</b>             | PAK1/PAK2/PAK3  |
| <b>Dilution:</b>              | Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications. |