



|  |                                    |  |
|--|------------------------------------|--|
|  | <h1>ARP-1 Polyclonal Antibody</h1> |  |
|--|------------------------------------|--|

|                                 |  |
|---------------------------------|--|
| <b>Swiss-Prot No.:</b>          | P24468   |
| <b>Molecular Weight:</b>        | 45.571   |
| <b>Applications:</b>            | WB,IHC,IF,ELISA  |
| <b>Reactivity:</b>              | Human,Mouse,Rat  |
| <b>Purification:</b>            | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  |
| <b>Immunogen:</b>               | Synthesized peptide derived from the N-terminal region of human ARP-1. at AA range: 1-80   |
| <b>Specificity/Sensitivity:</b> | ARP-1 Polyclonal Antibody detects endogenous levels of ARP-1 protein.  |
| <b>Other Names:</b>             | NR2F2; ARP1; TFCOUP2; COUP transcription factor 2; COUP-TF2; Apolipoprotein A-I regulatory protein 1; ARP-1; COUP transcription factor II; COUP-TF II; Nuclear receptor subfamily 2 group F member 2 |
| <b>Storage/Stability:</b>       | -20°C/1 year   |
| <b>Source:</b>                  | Rabbit   |
| <b>Form of Antibody:</b>        | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |
| <b>Protein Concentration:</b>   | 1 mg/ml  |

**For Research Use Only**

|                   |   |
|-------------------|---|
|                   |  <p>HuvEc</p> <p>(kD)</p> <p>117-</p> <p>85-</p> <p>48-</p> <p>34-</p> <p>26-</p> <p>19-</p> |
|                   | Western Blot analysis of various cells using ARP-1 Polyclonal Antibody  |
| <b>Gene Name:</b> | NR2F2   |
| <b>Dilution:</b>  | Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.                  |