




Btk Polyclonal Antibody

| | |
|---------------------------------|--|
| Swiss-Prot No.: | Q06187 |
| Molecular Weight: | 76.15 |
| Applications: | WB,IHC,ELISA |
| Reactivity: | Human,Mouse,Rat,Monkey |
| Purification: | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Immunogen: | Synthesized peptide derived from human Btk around the non-phosphorylation site of Y223. |
| Specificity/Sensitivity: | Btk Polyclonal Antibody detects endogenous levels of Btk protein. |
| Other Names: | BTK; AGMX1; ATK; BPK; Tyrosine-protein kinase BTK; Agammaglobulinaemia tyrosine kinase; ATK; B-cell progenitor kinase; BPK; Bruton tyrosine kinase |
| Storage/Stability: | -20°C/1 year |
| Source: | Rabbit |
| Form of Antibody: | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Protein Concentration: | 1 mg/ml |

For Research Use Only

| | |
|-------------------|--|
| | <p>MOUSE-BRAIN</p>  <p>170— 130— 100— 70— 55— 40— 35— 25— 15—</p> <p>BTK (Ab-222)</p> |
| | Western Blot analysis of various cells using Btk Polyclonal Antibody |
| Gene Name: | BTK |
| Dilution: | Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications. |