




## c-Fms Polyclonal Antibody

|                                 |   |
|---------------------------------|---|
| <b>Swiss-Prot No.:</b>          | P07333  |
| <b>Molecular Weight:</b>        | 107.984   |
| <b>Applications:</b>            | WB,ELISA  |
| <b>Reactivity:</b>              | Human   |
| <b>Purification:</b>            | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.                         |
| <b>Immunogen:</b>               | Synthesized peptide derived from human c-Fms around the non-phosphorylation site of Y699.   |
| <b>Specificity/Sensitivity:</b> | c-Fms Polyclonal Antibody detects endogenous levels of c-Fms protein.   |
| <b>Other Names:</b>             | CSF1R; FMS; Macrophage colony-stimulating factor 1 receptor; CSF-1 receptor; CSF-1-R; CSF-1R; M-CSF-R; Proto-oncogene c-Fms; CD antigen CD115 |
| <b>Storage/Stability:</b>       | -20°C/1 year  |
| <b>Source:</b>                  | Rabbit  |
| <b>Form of Antibody:</b>        | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   |
| <b>Protein Concentration:</b>   | 1 mg/ml   |

For Research Use Only

|                   |   |
|-------------------|---|
|                   |  <p>HepG2</p> <p>170—<br/>130—<br/>100—<br/>70—<br/>55—<br/>40—<br/>35—<br/>25—<br/>15—</p> <p>c-Fms</p> |
|                   | Western Blot analysis of HepG2 cells using c-Fms Polyclonal Antibody  |
| <b>Gene Name:</b> | CSF1R   |
| <b>Dilution:</b>  | Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.   |