




c-Src Polyclonal Antibody

| | |
|---------------------------------|---|
| Swiss-Prot No.: | P12931 |
| Molecular Weight: | 59.704 |
| Applications: | WB,IHC,IF,ELISA |
| Reactivity: | Human,Mouse,Rat |
| Purification: | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Immunogen: | Synthesized peptide derived from human c-Src around the non-phosphorylation site of Y419. |
| Specificity/Sensitivity: | c-Src Polyclonal Antibody detects endogenous levels of c-Src protein. |
| Other Names: | SRC; SRC1; Proto-oncogene tyrosine-protein kinase Src; Proto-oncogene c-Src; pp60c-src; p60-Src |
| Storage/Stability: | -20°C/1 year |
| Source: | Rabbit |
| Form of Antibody: | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Protein Concentration: | 1 mg/ml |

| | |
|-------------------|--|
| |  |
| | Western Blot analysis of various cells using c-Src Polyclonal Antibody |
| Gene Name: | SRC |
| Dilution: | Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications. |