

## CstF-64 Polyclonal Antibody

| Swiss-Prot No.:          | P33240  |
|--------------------------|---|
| Molecular Weight:        | 60.959  |
| Applications:            | WB,IHC,IF,ELISA   |
| Reactivity:              | Human, Mouse  |
| Purification:            | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.                       |
| Immunogen:               | Synthesized peptide derived from the N-terminal region of human CstF-64. at AA rangle: 30-110   |
| Specificity/Sensitivity: | CstF-64 Polyclonal Antibody detects endogenous levels of CstF-64 protein.   |
| Other Names:             | CSTF2; Cleavage stimulation factor subunit 2; CF-1 64 kDa subunit; Cleavage stimulation factor 64 kDa subunit; CSTF 64 kDa subunit; CstF-64 |
| Storage/Stability:       | -20°C/1 year  |
| Source:                  | Rabbit  |
| Form of Antibody:        | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   |
| Protein Concentration:   | 1 mg/ml   |

|            | 293   |
|------------|---|
|            | (kD)  |
|            | 117-  |
|            | 85-   |
|            |   |
|            |   |
|            | 48-   |
|            | 34-   |
|            |   |
|            | 26-   |
|            |   |
|            | 19-   |
|            |   |
|            | Western Blot analysis of various cells using CstF-64 Polyclonal |
|            | Antibody  |
| Gene Name: | CSTF2   |
|            |   |
|            | Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 -     |
| Dilution:  | 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA:               |
|            | 1/5000. Not yet tested in other applications.                   |
|            |   |