




## Cyclin G Polyclonal Antibody

<b>Swiss-Prot No.:</b>	P51959
<b>Molecular Weight:</b>	34.074
<b>Applications:</b>	WB,IHC,ELISA
<b>Reactivity:</b>	Human,Mouse,Rat
<b>Purification:</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Immunogen:</b>	Synthesized peptide derived from the Internal region of human Cyclin G. at AA range: 130-210
<b>Specificity/Sensitivity:</b>	Cyclin G Polyclonal Antibody detects endogenous levels of Cyclin G protein.
<b>Other Names:</b>	CCNG1; CCNG; CYCG1; Cyclin-G1; Cyclin-G
<b>Storage/Stability:</b>	-20°C/1 year
<b>Source:</b>	Rabbit
<b>Form of Antibody:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Protein Concentration:</b>	1 mg/ml

	<p style="text-align: center;">Jurkat</p>  <p>(kD)</p> <p>117-</p> <p>85-</p> <p>48-</p> <p>34-</p> <p>26-</p> <p>19-</p>
	Western Blot analysis of various cells using Cyclin G Polyclonal Antibody
Gene Name:	CCNG1
Dilution:	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications.