




## Cystatin C Polyclonal Antibody

<b>Swiss-Prot No.:</b>	P01034
<b>Molecular Weight:</b>	15.799
<b>Applications:</b>	WB,ELISA
<b>Reactivity:</b>	Human,Mouse,Rat,Monkey
<b>Purification:</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Immunogen:</b>	Synthesized peptide derived from the Internal region of human Cystatin C. at AA range: 60-140
<b>Specificity/Sensitivity:</b>	Cystatin C Polyclonal Antibody detects endogenous levels of Cystatin C protein.
<b>Other Names:</b>	CST3; Cystatin-C; Cystatin-3; Gamma-trace; Neuroendocrine basic polypeptide; Post-gamma-globulin
<b>Storage/Stability:</b>	-20°C/1 year
<b>Source:</b>	Rabbit
<b>Form of Antibody:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Protein Concentration:</b>	1 mg/ml

	
	Western Blot analysis of various cells using Cystatin C Polyclonal Antibody
Gene Name:	CST3
Dilution:	Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.