




## DCC Polyclonal Antibody

|                                 |  |
|---------------------------------|--|
| <b>Swiss-Prot No.:</b>          | P43146   |
| <b>Molecular Weight:</b>        | 158.457  |
| <b>Applications:</b>            | WB, IF, ELISA  |
| <b>Reactivity:</b>              | Human, Mouse, Rat  |
| <b>Purification:</b>            | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.                          |
| <b>Immunogen:</b>               | Synthesized peptide derived from the Internal region of human DCC. at AA rangle: 410-490   |
| <b>Specificity/Sensitivity:</b> | DCC Polyclonal Antibody detects endogenous levels of DCC protein.  |
| <b>Other Names:</b>             | DCC; IGDCC1; Netrin receptor DCC; Colorectal cancer suppressor; Immunoglobulin superfamily DCC subclass member 1; Tumor suppressor protein DCC |
| <b>Storage/Stability:</b>       | -20°C/1 year   |
| <b>Source:</b>                  | Rabbit   |
| <b>Form of Antibody:</b>        | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |
| <b>Protein Concentration:</b>   | 1 mg/ml  |

|            |   |
|------------|---|
|            |                                        |
|            | Western Blot analysis of 293 cells using DCC Polyclonal Antibody  |
| Gene Name: | DCC   |
| Dilution:  | Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications. |