




DOC-1 Polyclonal Antibody

| | |
|---------------------------------|--|
| Swiss-Prot No.: | 014519 |
| Molecular Weight: | 12.365 |
| Applications: | WB,ELISA |
| Reactivity: | Human,Mouse |
| Purification: | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Immunogen: | Synthesized peptide derived from the Internal region of human DOC-1. at AA range: 30-110 |
| Specificity/Sensitivity: | DOC-1 Polyclonal Antibody detects endogenous levels of DOC-1 protein. |
| Other Names: | CDK2AP1; CDKAP1; DOC1; Cyclin-dependent kinase 2-associated protein 1; CDK2-associated protein 1; Deleted in oral cancer 1; DOC-1; Putative oral cancer suppressor |
| Storage/Stability: | -20°C/1 year |
| Source: | Rabbit |
| Form of Antibody: | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Protein Concentration: | 1 mg/ml |

For Research Use Only

| | |
|------------|--|
| |  <p>HeLa</p> <p>(kD)</p> <p>117-</p> <p>85-</p> <p>48-</p> <p>34-</p> <p>26-</p> <p>19-</p> |
| | Western Blot analysis of various cells using DOC-1 Polyclonal Antibody |
| Gene Name: | CDK2AP1 |
| Dilution: | Western Blot: 1/500 - 1/2000. ELISA: 1/5000. Not yet tested in other applications. |