




## EMMPRIN Polyclonal Antibody

<b>Swiss-Prot No.:</b>	P35613
<b>Molecular Weight:</b>	42.2
<b>Applications:</b>	WB,IHC,ELISA
<b>Reactivity:</b>	Human,Mouse,Rat
<b>Purification:</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Immunogen:</b>	Synthesized peptide derived from the C-terminal region of human EMMPRIN. at AA range: 310-390
<b>Specificity/Sensitivity:</b>	EMMPRIN Polyclonal Antibody detects endogenous levels of EMMPRIN protein.
<b>Other Names:</b>	BSG; Basigin; 5F7; Collagenase stimulatory factor; Extracellular matrix metalloproteinase inducer; EMMPRIN; Leukocyte activation antigen M6; OK blood group antigen; Tumor cell-derived collagenase stimulatory factor; TCSF; CD antigen CD147
<b>Storage/Stability:</b>	-20°C/1 year
<b>Source:</b>	Rabbit
<b>Form of Antibody:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

**For Research Use Only**

<b>Protein Concentration:</b>	1 mg/ml
	<div> <div> <div>HUVEC</div> <div>RAW264.7</div> </div>  <div>CD147</div> </div>
	Western Blot analysis of various cells using EMMPRIN Polyclonal Antibody
<b>Gene Name:</b>	BSG
<b>Dilution:</b>	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications.