




## Endo180 Polyclonal Antibody

<b>Swiss-Prot No.:</b>	Q9UBG0
<b>Molecular Weight:</b>	166.655
<b>Applications:</b>	WB,IHC,IF,ELISA
<b>Reactivity:</b>	Human,Mouse,Rat
<b>Purification:</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Immunogen:</b>	Synthesized peptide derived from the N-terminal region of human Endo180. at AA rangle: 90-170
<b>Specificity/Sensitivity:</b>	Endo180 Polyclonal Antibody detects endogenous levels of Endo180 protein.
<b>Other Names:</b>	MRC2; CLEC13E; END0180; KIAA0709; UPARAP; C-type mannose receptor 2; C-type lectin domain family 13 member E; Endocytic receptor 180; Macrophage mannose receptor 2; Urokinase-type plasminogen activator receptor-associated protein; UPAR-asso
<b>Storage/Stability:</b>	-20°C/1 year
<b>Source:</b>	Rabbit
<b>Form of Antibody:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

<b>Protein Concentration:</b>	1 mg/ml
	
	Western Blot analysis of various cells using Endo180 Polyclonal Antibody
<b>Gene Name:</b>	MRC2
<b>Dilution:</b>	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.