




## HDAC2 Polyclonal Antibody

<b>Swiss-Prot No.:</b>	Q92769
<b>Molecular Weight:</b>	55.364
<b>Applications:</b>	WB,IHC,ELISA
<b>Reactivity:</b>	Human,Mouse,Rat,Monkey
<b>Purification:</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Immunogen:</b>	Synthesized peptide derived from human HDAC2 around the non-phosphorylation site of S394.
<b>Specificity/Sensitivity:</b>	HDAC2 Polyclonal Antibody detects endogenous levels of HDAC2 protein.
<b>Other Names:</b>	HDAC2; Histone deacetylase 2; HD2
<b>Storage/Stability:</b>	-20°C/1 year
<b>Source:</b>	Rabbit
<b>Form of Antibody:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Protein Concentration:</b>	1 mg/ml

For Research Use Only

	 <p>(kD)</p> <p>117- 85- 48- 34- 26- 19-</p> <p>Jurkat 3T3</p> <p>HDAC5</p>
	Western Blot analysis of various cells using HDAC2 Polyclonal Antibody
<b>Gene Name:</b>	HDAC2
<b>Dilution:</b>	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications.