




## JIK Polyclonal Antibody

<b>Swiss-Prot No.:</b>	Q9H2K8
<b>Molecular Weight:</b>	105.406
<b>Applications:</b>	WB,ELISA
<b>Reactivity:</b>	Human,Mouse,Rat
<b>Purification:</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Immunogen:</b>	Synthesized peptide derived from the Internal region of human JIK. at AA range: 500-580
<b>Specificity/Sensitivity:</b>	JIK Polyclonal Antibody detects endogenous levels of JIK protein.
<b>Other Names:</b>	TAOK3; DPK; JIK; KDS; MAP3K18; Serine/threonine-protein kinase TAO3; Cutaneous T-cell lymphoma-associated antigen HD-CL-09; CTCL-associated antigen HD-CL-09; Dendritic cell-derived protein kinase; JNK/SAPK-inhibitory kinase; Jun kinase-inhi
<b>Storage/Stability:</b>	-20°C/1 year
<b>Source:</b>	Rabbit
<b>Form of Antibody:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

**For Research Use Only**

<b>Protein Concentration:</b>	1 mg/ml
	 <p>KDa Jurkat 118 90 50 39 27 -JIK</p>
	Western Blot analysis of various cells using JIK Polyclonal Antibody
<b>Gene Name:</b>	TAOK3
<b>Dilution:</b>	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.