



|  |                                    |  |
|--|------------------------------------|--|
|  | <h1>LIR-7 Polyclonal Antibody</h1> |  |
|--|------------------------------------|--|

|                                 |  |
|---------------------------------|--|
| <b>Swiss-Prot No.:</b>          | Q8N149   |
| <b>Molecular Weight:</b>        | 52.992   |
| <b>Applications:</b>            | WB,IF,ELISA  |
| <b>Reactivity:</b>              | Human  |
| <b>Purification:</b>            | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  |
| <b>Immunogen:</b>               | Synthesized peptide derived from the Internal region of human LIR-7. at AA range: 20-100   |
| <b>Specificity/Sensitivity:</b> | LIR-7 Polyclonal Antibody detects endogenous levels of LIR-7 protein.  |
| <b>Other Names:</b>             | LILRA2; ILT1; LIR7; Leukocyte immunoglobulin-like receptor subfamily A member 2; CD85 antigen-like family member H; Immunoglobulin-like transcript 1; ILT-1; Leukocyte immunoglobulin-like receptor 7; LIR-7; CD antigen CD85h |
| <b>Storage/Stability:</b>       | -20°C/1 year   |
| <b>Source:</b>                  | Rabbit   |
| <b>Form of Antibody:</b>        | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |
| <b>Protein Concentration:</b>   | 1 mg/ml  |

**For Research Use Only**

|                   |   |
|-------------------|---|
|                   |                                        |
|                   | Western Blot analysis of various cells using LILRA2 Polyclonal Antibody   |
| <b>Gene Name:</b> | LILRA2  |
| <b>Dilution:</b>  | Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications. |