

Lunatic Fringe Polyclonal Antibody

| Swiss-Prot No.: | Q8NES3 | | |
|--------------------------|---|--|--|
| Molecular Weight: | 41.773 | | |
| Applications: | WB,ELISA | | |
| Reactivity: | Human,Mouse,Rat | | |
| Purification: | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. | | |
| Immunogen: | Synthesized peptide derived from the Internal region of human Lunatic Fringe. at AA rangle: 90-170 | | |
| Specificity/Sensitivity: | Lunatic Fringe Polyclonal Antibody detects endogenous levels of Lunatic Fringe protein. | | |
| Other Names: | LFNG; Beta-1; 3-N-acetylglucosaminyltransferase lunatic fringe; O-fucosylpeptide 3-beta-N-acetylglucosaminyltransferase | | |
| Storage/Stability: | -20°C/1 year | | |
| Source: | Rabbit | | |
| Form of Antibody: | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. | | |
| Protein Concentration: | 1 mg/ml | | |

| | 178 100 70 55 40 35 25 | 3T3 | Lunatic Fringe | |
|------------|---|-----|----------------|--|
| | Western Blot analysis of 3T3 cells using Lunatic Fringe Polyclonal Antibody | | | |
| Gene Name: | LFNG | | | |
| Dilution: | Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. | | | |