




## MDM2 Polyclonal Antibody

<b>Swiss-Prot No.:</b>	Q00987
<b>Molecular Weight:</b>	55.233
<b>Applications:</b>	WB,IHC,IF,ELISA
<b>Reactivity:</b>	Human,Mouse,Monkey
<b>Purification:</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Immunogen:</b>	Synthesized peptide derived from human MDM2 around the non-phosphorylation site of S166.
<b>Specificity/Sensitivity:</b>	MDM2 Polyclonal Antibody detects endogenous levels of MDM2 protein.
<b>Other Names:</b>	MDM2; E3 ubiquitin-protein ligase Mdm2; Double minute 2 protein; Hdm2; Oncoprotein Mdm2; p53-binding protein Mdm2
<b>Storage/Stability:</b>	-20°C/1 year
<b>Source:</b>	Rabbit
<b>Form of Antibody:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Protein Concentration:</b>	1 mg/ml

For Research Use Only

	 <p>SH-SY5Y COLO</p> <p>170— 130— 100— 70— 55— 45— 35— 25— 15—</p> <p>MDM2</p>
	Western Blot analysis of various cells using MDM2 Polyclonal Antibody
<b>Gene Name:</b>	MDM2
<b>Dilution:</b>	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications.