




Met Polyclonal Antibody

Swiss-Prot No.:	P08581
Molecular Weight:	155.527
Applications:	WB,IF,ELISA
Reactivity:	Human,Mouse,Rat
Purification:	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Immunogen:	Synthesized peptide derived from human Met around the non-phosphorylation site of Y1003.
Specificity/Sensitivity:	Met Polyclonal Antibody detects endogenous levels of Met protein.
Other Names:	MET; Hepatocyte growth factor receptor; HGF receptor; HGF/SF receptor; Proto-oncogene c-Met; Scatter factor receptor; SF receptor; Tyrosine-protein kinase Met
Storage/Stability:	-20°C/1 year
Source:	Rabbit
Form of Antibody:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Protein Concentration:	1 mg/ml

For Research Use Only

	 Western blot analysis of HepG2 cells using Met Polyclonal Antibody. The blot shows a single band at approximately 145 kD. Molecular weight markers are indicated on the left at 170, 130, 95, 72, and 55 kD.
	Western Blot analysis of various cells using Met Polyclonal Antibody
Gene Name:	MET
Dilution:	Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications.